

2360 gulf freeway south, Suite 106, League-city, TX 77573. Ph: 281-337-3222. Fax 281-337-9222

SEALANT CONSENT:	Tooth #	
I, measure intended to facil the teeth.	, understand that the treatment of teeth through the use of silitate the inhibition of dental caries (tooth decay or cavities) in the pits and fissures of the	
	risks, if any, which may be associated with the placement of sealants even thoug . Mbibi, and/or a sealant certified dental staff at Tot's N Teens Pediatric Dental Spe	
In addition to possible ufollowing:	insuccessful results and failure of the sealant, the risks of the procedure include but	are not limited to the
etches the surface en caustic and if the pat amount of the solute	Teeth: The teeth are prepared through use of an enamel etching technique: using a spectramel in the area in which the sealant is to be placed to aid in its retention. The etching tient makes any unexpected movements during the application process there is the partion may attach to the soft tissues of the mouth which could cause some slight is a possibility. If the etching solution contacts the root surface, the tooth may developed the soft tissues of the mouth which could cause some slight in a possibility.	g solution is somewhat possibility that <b>a small</b> <b>ht tissue burns</b> . This
length of time over which including, but not limited to a. The forces of patient than in an b. The types of four sticky candies a sealant.  c. Inadequate or sealant causing as Entire tooth is not proof the teeth. These pits at flow into and seal those	slodging of the sealant: There is the possibility of the sealant loosening or becoming dish this may happen is indeterminable because of the many variables which can impact to the following:  If mastication (chewing). These forces differ from patient to patient. The forces may be another. Also, the way teeth occlude (come together in chewing) may have an effect on the food or other substances that are put in the mouth and chewed. Very sticky foods such a such as caramels; some licorices; very hard substances, etc; may cause loosening ral hygiene such as infrequent or improper brushing of the teeth also may allow leakage it to loosen and allow decay to develop.  Totected with sealants: Sealants are applied primarily to the pits and fissures that are in and fissures are extremely susceptible to decay and can be protected through the application areas. However, sealants do not protect the areas between the teeth, so thorough brush is is necessary. Otherwise, decay could develop in those areas uncovered by the sealants.	the life of the sealant e much greater in one le life of the sealants. as some types of gum; or dislodgment of the around and under the attention of sealants which ushing and the use of
	understand that it is my responsibility to notify to ccur or if any problems relating to the treatment rendered are experienced. Rou ended to allow ongoing assessment of the sealants placed.	
received answers to my a rendered though no guara of substantial harm, if a	I have been given the opportunity to ask questions regarding the nature and purpose satisfaction. I voluntarily undergo this treatment in hopes of achieving the desired resurantees have been made regarding the outcome. I hereby assume any and all possible rany, which may be associated with any phase of this treatment. The fee(s) for these cept them as satisfactory. By signing this form, I am freely giving my consent to authorize	ults from the treatment risks, including the risk e services have been
	and/or all associates involved in rendering the services or treatment necessary to the ex on and/or prescribing of any anesthetic agents and/or medications.	isting dental condition,
	Patient's name (please print)	Date
	Signature of legal guardian	Witness Signature